

Potential role of *Thermus thermophilus* and *T. oshimai* in high rates of nitrous oxide (N₂O) production in ~80 °C hot springs in the US Great Basin

B. P. HEDLUND,¹ A. I. MCDONALD,¹ J. LAM,¹ J. A. DODSWORTH,¹ J. R. BROWN² AND B. A. HUNGATE²

¹School of Life Sciences, University of Nevada, Las Vegas, Las Vegas, NV, USA

²Department of Biological Sciences and Merriam-Powell Center for Environmental Research, Northern Arizona University, Flagstaff, AZ, USA

SUMMARY

Ambient nitrous oxide (N₂O) emissions from Great Boiling Spring (GBS) in the US Great Basin depended on temperature, with the highest flux, $67.8 \pm 2.6 \mu\text{mol N}_2\text{O-N m}^{-2} \text{ day}^{-1}$, occurring in the large source pool at 82 °C. This rate of N₂O production contrasted with negligible production from nearby soils and was similar to rates from soils and sediments impacted with agricultural fertilizers. To investigate the source of N₂O, a variety of approaches were used to enrich and isolate heterotrophic micro-organisms, and isolates were screened for nitrate reduction ability. Nitrate-respiring isolates were identified by 16S rRNA gene sequencing as *Thermus thermophilus* (31 isolates) and *T. oshimai* (three isolates). All isolates reduced nitrate to N₂O but not to dinitrogen and were unable to grow with N₂O as a terminal electron acceptor. Representative *T. thermophilus* and *T. oshimai* strains contained genes with 96–98% and 93% DNA identity, respectively, to the nitrate reductase catalytic subunit gene (*narG*) of *T. thermophilus* HB8. These data implicate *T. thermophilus* and *T. oshimai* in high flux of N₂O in GBS and raise questions about the genetic basis of the incomplete denitrification pathway in these organisms and on the fate of biogenic N₂O in geothermal environments.

Received 21 April 2011; accepted 22 August 2011

Corresponding author: Brian P. Hedlund. Tel.: (702) 895 0809; fax: (702) 895 3950; e-mail: brian.hedlund@unlv.edu

INTRODUCTION

Nitrous oxide (N₂O) is a powerful greenhouse gas that has become a major focus of climate change research in recent decades. Measurements of N₂O flux across soil/air and water/air interfaces from a wide variety of terrestrial and marine environments have documented dramatic increases in emissions owing to anthropogenic sources of fixed N, more than 70% of which can be linked with the use of agricultural fertilizers (Seitzinger *et al.*, 2000, 2006; EPA, 2010). In soils and both freshwater and coastal marine sediments, N₂O emissions can be elevated several orders of magnitude owing to the direct application of fertilizers or runoff (Matson *et al.*, 1998). Biological nitrification and denitrification are the two major sources of N₂O, with denitrification likely to be the dominant source (Zumft & Kroneck, 2007). N₂O is an intermediate in typical denitrification pathways in which dinitrogen

gas (N₂) is the terminal product. N₂O typically constitutes a trace emission from denitrifying microbial communities (e.g., <5% of N₂O + N₂), and globally N₂O is estimated to comprise ~2.6–3.9% of the total denitrification flux (Seitzinger, 1988; Schlesinger, 2009).

A wide diversity of bacteria and archaea (>60 genera) and even some eukaryotes are capable of denitrification, the vast majority of which catalyze complete denitrification with N₂ as the final product (Zumft & Kroneck, 2007). Micro-organisms with truncated denitrification pathways that terminate with N₂O have been known for several decades (Hart *et al.*, 1965), and genomics has revealed that this phenotype can result from mutations in the N₂O reductase (*nos*) genes or from the complete absence of *nos* genes (Zumft & Kroneck, 2007). In geothermal environments, a wide variety of thermophiles can respire nitrate or nitrite, including both *Crenarchaeota* (Völkl *et al.*, 1993; Blöchl *et al.*, 1997; Afshar *et al.*, 1998) and

Euryarchaeota (Hafenbradl *et al.*, 1996; Vorholt *et al.*, 1997) and members of several phyla of *bacteria*, such as the *Aquificae* (Eder & Huber, 2002; Gotz *et al.*, 2002; Takai *et al.*, 2003), *Firmicutes* (Huber *et al.*, 1996; Mishima *et al.*, 2009; Poli *et al.*, 2009), and *Thermales*. The *Thermales* include some of the best-studied nitrate- and nitrite-reducing thermophiles. Some isolates of *Thermus thermophilus* are capable of denitrification to N₂ (Cava *et al.*, 2008); however, most nitrate-reducing *Thermales* can only reduce nitrate to nitrite, including some isolates of *T. thermophilus* (Cava *et al.*, 2008) and all facultative anaerobic species of *Meiothermus*, *Oceanithermus*, and *Vulcanithermus* (Miroshnichenko *et al.*, 2003a,b; Pires *et al.*, 2005; Albuquerque *et al.*, 2009, 2010; Zhang *et al.*, 2010). Some strains of *T. oshimai* and *T. brockianus* are reported denitrifiers, but the denitrification phenotype has not been described in detail (da Costa *et al.*, 2001).

Considerable research has been carried out to elucidate the genetic and biochemical basis of nitrate reduction pathways in *T. thermophilus*. Genes encoding the ability to reduce nitrate to nitrite in *T. thermophilus* HB8 and NAR1 comprise three adjacent operons, *nar*, *nrc*, and *dnr* that are located on a megaplasmid, termed the nitrate conjugative element (NCE). The NCE megaplasmid can be transferred among *T. thermophilus* strains by conjugation (Ramirez-Arcos *et al.*, 1998). The *nar* operon encodes the membrane-bound nitrate reductase (*narG,H,I*), among other genes necessary for nitrate respiration (Ramirez *et al.*, 2000; Zafra *et al.*, 2002). The nitrate reductase alpha subunit, *narG*, is the highly conserved catalytic subunit and is commonly used as a functional biomarker for dissimilatory nitrate-reducing bacteria in molecular ecology studies (Philippot *et al.*, 2002).

Despite a strong foundation based on a large body of research in nonthermal environments and biochemistry of nitrate reduction pathways in *T. thermophilus*, very few studies have addressed denitrification in terrestrial geothermal environments (Burr *et al.*, 2005). A recent study reported coordinated measurements of gross ammonia oxidation, denitrification, dissimilatory nitrate reduction to ammonium (DNRA), microbial community composition, and abundance and distribution of ammonia monooxygenase large subunit genes (*amoA*) and *narG* in two ~80 °C springs in the US Great Basin, Great Boiling Spring (GBS) and Sandy's Spring West (SSW) (Dodsworth *et al.*, 2011). The study revealed an extremely active N-cycle, including abundant populations of ammonia-oxidizing archaea related to '*Ca. Nitrosocaldus yellowstonii*' that are likely responsible for the oxidation of source water ammonia to nitrite at rates comparable to those in nonthermal aquatic sediments. Denitrification rates were high and could be stimulated ~10-fold by the addition of nitrate and a complex organic mixture, implying an important role of heterotrophic denitrifiers, whereas possible electron donors for chemolithotrophy failed to stimulate denitrification. However, the identities

and activities of heterotrophic denitrifiers in the springs remained speculative.

Here, we report a substantial flux of N₂O from GBS and describe the isolation and characterization of a large number of heterotrophic nitrate-reducing micro-organisms from GBS and SSW. We show that all isolates produce N₂O as the terminal nitrate reduction product, regardless of sampling date, location, or isolation strategy, which is surprising in light of the absence of previous reports of any member of the *Thermales* with this phenotype. The potential implications of these results on the fate of microbially produced N₂O in terrestrial geothermal ecosystems are discussed.

MATERIALS AND METHODS

Sample sites, field chemistry, and measurement of *in situ* gas flux

The aqueous chemistry, mineralogy, and sediment and planktonic microbial communities inhabiting source pools of GBS and SSW have been described in detail previously (Costa *et al.*, 2009; Dodsworth *et al.*, 2011). Physical and limited chemical data collected on four additional sampling trips are summarized in Table 1. In the field, nitrate and nitrite concentrations were measured by diazotization with sulfanilamide and coupling with *N*-(1-naphthyl)-ethylenediamine dihydrochloride with or without cadmium reduction, respectively (LaMotte, Chestertown, MD, USA). Total ammonia was measured by the nesslerization method using a commercially available kit (LaMotte). For the measurement of dissolved N₂O, triplicate 40 mL water samples were collected in 160-mL serum bottles, cooled to ambient temperature (~25 °C), and shaken three times for 30 s at intervals of 3 min to equilibrate N₂O with the headspace. Gas samples were stored, transported, and analyzed for N₂O by GC-ECD as described in detail below for the measurements of N₂O flux.

On April 7–9, 2009, net fluxes of N₂O, methane (CH₄), and carbon dioxide (CO₂) at the air–water interface were measured using floating static chambers (MacIntyre *et al.*, 1995). In GBS, gas fluxes were measured at three locations, the center of the large source pool (Fig. 1; 82 °C), a shallow shelf

Table 1 Spring temperature, pH, and bulk water inorganic N chemistry (μM)

Site/date	Temp. (°C)	pH	NH ₄ ⁺	NO ₂ ⁻	NO ₃ ⁻
GBS					
Oct. 2008	81	7.20	39	4.2	16
Dec. 2008	83	7.15	70	0.79	1.8
April 2009	82	6.90	28	10.2	7.7
June 2009	84.4	6.85	92	1.0	8.8
SSW					
April 2009	79	6.73	43	0.86	<0.1

NH₄⁺, ammonia/ammonium; NO₂⁻, nitrite; NO₃⁻, nitrate.




Fig. 1 Photograph of GBS showing sites for measurement of *in situ* gas flux and sampling for microbial enrichments. Gas flux was measured in the center of the spring and sites B and E. Gas flux was also measured in grass-covered soil a few meters from the spring to the right and behind the photographer. Samples for microbial enrichments were collected on the shallow shelf at site B and a neuston biofilm at site C. A foil-covered tube rack with denitrification experiments described by Dodsworth *et al.* (2011) is visible at site B. GBS, Great Boiling Spring.

used for denitrification and DNRA field measurements (Fig. 1 Site B; 82 °C (Dodsworth *et al.*, 2011), and a shallow shelf hosting a thin cyanobacterial mat (Fig. 1 Site E; ~65 °C). Additional samples were collected in a patch of ambient-temperature soil hosting grasses a few meters from GBS. In SSW, samples were collected from the center of the source pool.

The static chambers were 1-cm walled PVC tubes, 30 cm diam. and 15 cm height, capped with a circular 1-cm-thick piece of flat PVC. A sample port was located on the top of each chamber and consisted of a stainless steel quick-connect bulkhead valve. A vinyl vent tube (1.3 cm diam. by 15 cm long) passed through the top of the chamber and was attached to the inside wall. The length and diameter of vent tube were calculated to reduce mixing of chamber air with outside air owing to perturbations from wind and sample collection. To measure gas fluxes, chambers were placed in inflatable inner tubes floating on the spring surface, such that the PVC chamber wall descended 3 cm into the water. Immediately after deploying the chamber and after 20 and 40 min, air samples were collected in 100 mL Silonite™ coated stainless steel sample canisters (Entech Instruments, Inc., Simi Valley, CA, USA) equipped with quick disconnect valves and evacuated prior to field sampling. Samples were taken by attaching one end of a sample canister to the chamber bulkhead fitting, such that the pre-evacuated canister immediately collected 100 mL of chamber air. To detect leaks during transport to the laboratory, the canister was then overpressurized as follows: A syringe was attached to the other end of the canister and an additional 60 mL of chamber air withdrawn. The syringe and canister were then together disconnected from the chamber, and the additional 60 mL was injected into the canister from the syringe. Canisters were transported back to the laboratory for the analysis of CO₂, CH₄, and N₂O concentrations. Ten

millilitres of gas was removed from each canister by syringe and injected into an Agilent Technologies Model 6890A gas chromatograph (Agilent Technologies, Inc., Palo Alto, California, USA). CO₂ and CH₄ concentrations were measured in He carrier using a flame ionization detector with in-line methanizer, and N₂O was measured using an electron capture detector, with an argon/methane carrier. Gas flux rates were calculated using linear regression of gas concentration over time and (accounting for chamber volume and area) expressed as moles of gas per m² per day.

Microbial enrichment and isolation

Information on the identity, source material, and enrichment strategy for each isolate is summarized in Table 2. All samples for enrichments were collected from water or surface sediments (top 1 cm) from a shallow shelf in GBS (Fig. 1; Site B) or a neuston biofilm collected in the outflow channel (Fig. 1; Site C). Samples from SSW were taken from shallows in the small source pool about 5–10 cm below the water–air interface at the east side of the spring pool at the site described (Dodsworth *et al.*, 2011).

Four different enrichment strategies were used, as follows. For strategy 1, sediment was collected in sterile polypropylene tubes and transported to the laboratory and stored in the dark without redox or temperature control. Within a week of sampling, samples were used to inoculate 40 mL of aerobic or anaerobic (N₂-sparged; 2 atm N₂) GBS spring water ultrafiltrate amended with 0.1% peptone and 0.05% yeast extract in stoppered serum bottles (e.g., Wheaton 160 mL bottles, 223748). Anaerobic enrichments were supplemented with 9 mM NaNO₃. For strategy 2, spring water or a dilute spring water/sediment slurry was transferred into a serum bottle, fortified with 0.1% peptone and 0.05% yeast extract, and immediately incubated under aluminum foil in the shallows of the spring without allowing the sample to cool. For strategy 3, Castenholz medium D, which contains 9 mM NO₃⁻ (Castenholz, 1969), in borosilicate screw-capped tubes (3 mL per tube) was amended with a variety of carbon substrates, inoculated with hot, freshly collected samples, and incubated with an air headspace in the shallows of the springs under aluminum foil. Successful enrichments used for microbial isolation shown in Table 2 include lactate (1 mM), an equimolar mixture of formate, lactate, acetate, and propionate (1 mM each), BSA (0.01%), yeast extract and peptone (0.05%), and casamino acids (0.05%). For strategy 4, sediment slurries were transported to the laboratory without temperature or redox control and inoculated into the medium of Allen (Allen, 1959), which was modified to pH 7 and by the addition of 0.01% yeast extract and 20 mM sucrose or 0.5% potato starch. Turbid cultures were plated onto the same medium solidified with gelrite.

All enrichments were incubated for 12–36 h, and contents were plated onto Castenholz medium D supplemented with

Table 2 Summary of source material and enrichment strategy for each nitrate-reducing isolate

Strain*	Source	Enrichment/isolation (strategy†)
<i>Thermus thermophilus</i>		
SSW-01	SSW sediment (10/2008)	Lab, aerobic, YE/Pept, 75 °C (1)
SSW-06 (JL-6)*	SSW sediment (12/2008)	Field, aerobic, YE/Pept, 75 °C (2)
SSW-10	SSW sediment (12/2008)	Field, aerobic, YE/Pept, 75 °C (2)
SSW-11	SSW sediment (12/2008)	Field, aerobic, YE/Pept, 75 °C (2)
SSW-14	SSW sediment (12/2008)	Field, aerobic, YE/Pept, 75 °C (2)
SSW-18b (JL-18)*	SSW water (12/2008)	Field, aerobic, YE/Pept, 75 °C (2)
SSW-19	SSW sediment (12/2008)	Field, aerobic, YE/Pept, 75 °C (2)
SSW-20	SSW sediment (12/2008)	Field, aerobic, YE/Pept, 75 °C (2)
SSW-21	SSW sediment (12/2008)	Field, aerobic, YE/Pept, 75 °C (2)
GBS-02	GBS sediment (10/2008)	Laboratory, aerobic, YE/Pept, 75 °C (1)
GBS-03	GBS water (10/2008)	Field, aerobic, YE/Pept, 75 °C (2)
GBS-04	GBS water (10/2008)	Field, aerobic, YE/Pept, 75 °C (2)
GBS-05	GBS water (10/2008)	Field, aerobic, YE/Pept, 75 °C (2)
GBS-08	GBS sediment (12/2008)	Field, aerobic, YE/Pept, 75 °C (2)
GBS-09	GBS sediment (12/2008)	Field, aerobic, YE/Pept, 75 °C (2)
GBS-10	GBS sediment (12/2008)	Field, aerobic, YE/Pept, 75 °C (2)
GBS-12	GBS sediment (12/2008)	Field, aerobic, YE/Pept, 75 °C (2)
GBS-27	GBS sediment (06/2009)	Field, aerobic, YE/Pept, 70 °C (3)
GBS-32	GBS sediment (06/2009)	Field, anaerobic, YE/Pept, 70 °C (3)
GBS-33	GBS sediment (06/2009)	Field, aerobic, YE/Pept, 70 °C (3)
GBS-34	GBS sediment (06/2009)	Field, aerobic, Casamino acids, 70 °C (3)
GBS-36	GBS sediment (06/2009)	Field, aerobic, Casamino acids, 70 °C (3)
GBS-40	GBS neuston (06/2009)	Field, aerobic, Lactate, 70 °C (3)
GBS-44	GBS sediment (06/2009)	Field, aerobic, YE/Pept, 70 °C (3)
GBS-45	GBS sediment (06/2009)	Field, aerobic, BSA, 70 °C (3)
GBS-49	GBS sediment (06/2009)	Field, aerobic, FLAP, 70 °C (3)
GBSBH-03	GBS sediment (12/2008)	Laboratory, anaerobic, YE/Pept, 75 °C (1)
GBSBH-05	GBS sediment (12/2008)	Laboratory, anaerobic, YE/Pept, 75 °C (1)
GBSBH-09	GBS sediment (12/2008)	Laboratory, anaerobic, YE/Pept, 75 °C (1)
PS1	GBS sediment (3/08)	Laboratory, aerobic, starch, 76 °C (4)
S1	GBS sediment (3/08)	Laboratory, aerobic, sucrose, 76 °C (4)
<i>T. oshimai</i>		
SSW-02 (JL-2)*	SSW sediment (10/2008)	Laboratory, aerobic, YE/Pept, 75 °C (1)
SSW-03	SSW sediment (12/2008)	Field, aerobic, YE/Pept, 75 °C (2)
SSW-04 (JL-4)*	SSW sediment (12/2008)	Field, aerobic, YE/Pept, 75 °C (2)

GBS, great boiling spring.

*Strains used for detailed nitrate reduction stoichiometry experiments. Isolate synonyms in parenthesis are used in Fig. 2.

†The four enrichment strategies are described in detail in the Materials and Methods. Carbon substrate abbreviations are as follows: YE/Pept, yeast extract, and peptone (0.05% each); BSA, bovine serum albumin (0.01%); FLAP, formate, lactate, acetate, and propionate (1 mM each).

0.1% yeast extract and 0.1% tryptone (Castenholz, 1969) solidified with 2% agar or modified Allen medium (strategy 4 only, described earlier). Anaerobic cultures were plated in a Coy Type B anaerobic chamber (Coy Laboratory Products Inc., Grass Lake, MI, USA) containing an atmosphere of N₂ (~90%), CO₂ (~5%), and H₂ (~5%) and incubated under an atmosphere of N₂ (2 atm) in aluminum pressure cups (model 51-600; C.A. Technologies, Boulder, CO, USA) sealed with viton o-rings (#9464K673; McMaster Carr, Robbinsville, NJ, USA). Pressure cups were modified for use as anaerobic growth chambers by sealing all ports with high temperature gasket material (#1967; NEA products Co., Belleville, NJ, USA) and silicone sealant (#81160; Permatex, Solon, OH, USA). For all enrichments, several colonies representing each colony morphology type were picked and streaked onto the

same medium until axenic cultures were obtained (3–5× streak plates). Pure cultures were immediately used to make –80 °C freezer stocks in media supplemented with 15% glycerol. Each strain was revived from the freezer stocks, streaked to check purity, and used without delay for nitrate reduction experiments described below.

DNA isolation, 16S rRNA and narG PCR and DNA sequencing

DNA was extracted from isolates using a colony lysis buffer (Johnson *et al.*, 2001), and 16S rRNA genes were PCR-amplified using primers specific for bacteria: 9bF (Eder *et al.*, 1999) and 1512uR (Eder *et al.*, 2001). The 25 µL PCR mixture contained 1 µL DNA extract, 200 nM of each primer,

200 µM each dNTP (Promega, Madison, WI, USA), 0.65 U of GoTaq DNA polymerase (Promega), and 5 µL of 5× GoTaq buffer (Promega). Cycling conditions were denaturation at 96 °C for 4 min followed by 35 cycles of denaturation (30 s at 94 °C), primer annealing (30 s at 55 °C), and elongation (1.5 min at 72 °C), with a final elongation step (10 min at 72 °C). PCR for *narG* was carried out using the primers narG1960f and narG2650r and using the reagents described above and the cycling conditions described by Phillipot (Phillipot *et al.*, 2002), except that the final extension was shortened to 10 min. For both 16S rRNA genes and *narG*, PCR products were sequenced using the Sanger method at Functional Biosciences, Madison, WI, using the forward PCR primer. All sequences had >700 base pairs with Phred score >20. Sequences were identified using BLASTn (Altschul *et al.*, 1990). A section of *narG* corresponding to the *T. thermophilus* HB8 amino acid position 581–753 was aligned with related *narG* and deduced NarG sequences using default parameters of ClustalW within BioEdit (Hall, 2005). All alignments were checked manually. 16S rRNA and *narG* sequences have been deposited to GenBank with accession numbers JN115057–JN115090 and JN115091–JN115096, respectively.

Screen for nitrate reduction phenotype

Isolates were initially screened for nitrate reduction phenotype by testing the ability to grow in anaerobic Castenholz medium D with 0.1% yeast extract and 0.1% tryptone under N₂ with 9 mM nitrate as the sole terminal electron acceptor in Balch tubes (Bellco 2048-00150) sealed with butyl rubber stoppers (Bellco 2048-18150) and aluminum seals (Bellco 2048-11020). The ability to produce N₂O was determined by GC-ECD analysis of headspace gas after 72 h, as described in detail below. The ability to reduce N₂O to N₂ was tested in the same medium under a 100% He atmosphere by testing their ability to produce N₂ as assessed by production of gas in Durham vials and by GC-TCD, as described in detail below. In addition, late-exponential phase (72 h) cells grown anaerobically with 9 mM nitrate were tested for their ability to grow or produce gas on the same medium but with 100% N₂O (2 atm) as the sole terminal electron acceptor.

Measurement of cell density, nitrate, and denitrification products during anaerobic growth

For all nitrate reduction experiments, isolates were grown in Castenholz medium D with 0.1% yeast extract and 0.1% tryptone that was sparged for 45 min with N₂ and dispensed in the anaerobic chamber described earlier. The headspace was changed prior to autoclaving by 10 cycles of evacuation (1 min) and filling to 2 atm with 99.999% helium.

Cultures were revived from freezer stocks by plating onto Castenholz medium D with 0.1% yeast extract and 0.1% tryptone.

Colonies were resuspended in small volumes of anaerobic medium and transferred to serum bottles containing a 100% helium atmosphere (2 atm) prepared as described earlier. Cells were grown and passed using He-rinsed syringes and a 1:50 inoculum into the same prewarmed medium twice prior to the experiment to dilute contaminating N₂ and ensure nitrate reduction pathways were active. Cultures were incubated at 75 °C with rotary shaking at 100 rpm with serum bottles secured in a horizontal position to maximize gas equilibration.

Aqueous medium (0.3 mL) and headspace gas (2.5 mL) were collected at each time point using He-flushed syringes for the measurement of cell density, aqueous nitrate, nitrite, and ammonia concentrations, and concentrations of headspace N₂O and N₂. Cells were counted using a Petroff-Hausser chamber and an Olympus BX-51 phase-contrast microscope with brightness and contrast optimized by using PICTUREFRAME software (Optronics, Goleta, CA, USA). Nitrate concentration was measured by nitration of 2-hydroxybenzoic acid at acidic pH, which does not suffer from nitrite interference (Bhandari & Simlot, 1986). Nitrite and total ammonia concentrations were measured by miniaturization of commercially available kits using methods described above for field measurements (LaMotte). Headspace N₂O concentration was determined by GC-ECD by the injection of a 2.0-mL sample into a GC-2014 Nitrous Oxide Analyzer (Shimadzu, Moorpark, CA, USA), modified and operated as described (Dodsworth *et al.*, 2010). At N₂O concentrations >100 ppm, a nonlinear equation was applied to generate and optimize a standard curve using the open-source curve fitting tools hosted at zunzun.com. Headspace N₂ concentration was determined by GC-TCD by the injection of a 0.5-mL sample into a GC-2014 gas chromatograph (Shimadzu) using 99.999% argon at 40 mL min⁻¹ as the carrier gas and a Supelco 80/100 mesh Molecular Sieve column (Sigma-Aldrich, St. Louis, MO, USA). Injector, column, and detector temperatures were 150, 80, and 210 °C, respectively. For both gases, headspace concentrations were used to calculate aqueous gas concentrations and subsequently the total molar amount of each gas by using Henry's Law and constants modified for high temperature (Wilhelm *et al.*, 1977).

RESULTS AND DISCUSSION

In situ N chemistry and gas flux

Consistent with the previous reports (Costa *et al.*, 2009; Dodsworth *et al.*, 2011), ammonia concentrations were higher than concentrations of nitrate plus nitrite in GBS and SSW at all sampling dates, particularly in SSW (Table 1). This is consistent with a model in which the common subterranean reservoir that sources these springs is reduced, with essentially no nitrate or nitrite. Because SSW has a short water residence time (5–15 min.), oxidized products of chemolithotrophic




Fig. 2 Gas flux measured in April 2009 at GBS the center of the spring (main pool; 82 °C), site B (82 °C), site E (~65 °C), adjacent grass-covered soil, and in the main pool of SSW. Bars represent the mean and standard error ($n = 6-7$ for GBS and soil locations; $n = 2$ for SSW). Letter designations above bars denote levels that are significantly different from each other using one-way ANOVA. GBS, Great Boiling Spring; SSW, Sandy's Spring West.

ammonia oxidation do not accumulate in bulk spring water. In contrast, the higher levels of nitrate and nitrite in GBS are owing to a much longer water residence time (>1 day) and consistent with the high rates of ammonia oxidation recently reported (Dodsworth *et al.*, 2011).

N₂O flux across the water/atmosphere interface measured in April 2009 was highest at the center of the GBS source pool ($67.8 \pm 2.6 \mu\text{mol N m}^{-2} \text{ day}^{-1}$; 82 °C) and lower but significant on a shallow shelf (Fig. 1, B site; $15.3 \pm 2.5 \mu\text{mol N m}^{-2} \text{ day}^{-1}$; 82 °C; Fig. 2). In contrast, N₂O flux from a submerged, thin cyanobacterial mat (Fig. 2 E site; ~65 °C), nearby grass-covered soil, and the SSW source pool were low or negligible. This pattern of N₂O production contrasted with fluxes of CO₂ and CH₄, which were highest in the SSW source pool (Fig. 2). CO₂ has been suggested to be geogenic in some springs in the GBS system based on relatively high $\delta^{13}\text{C}$ values of free-phase CO₂ and carbonate solids [~ -6 per mil (Romanek *et al.*, 2005)]. The relative enrichment of N₂O in the GBS source pool and shallow shelf, when compared with geogenic CO₂ (which were approximately equal in GBS and SSW), is consistent with a biogenic source of N₂O within the source pool and in shallow, high temperature sediments.

Nitrification and denitrification were highly active in GBS in October 2008 and April 2009, so both nitrification and

denitrification are possible sources of biogenic N₂O in GBS (Dodsworth *et al.*, 2011). Acetylene block experiments with sediment slurries revealed denitrification rates at the GBS B site of $2,352 \pm 240 \mu\text{mol N m}^{-2} \text{ day}^{-1}$ in October 2008 and $10,422 \pm 92.2 \mu\text{mol N m}^{-2} \text{ day}^{-1}$ in April 2009. Similar unblocked experiments suggested a rate of N₂O flux of approximately an order of magnitude lower. Although sediment slurry experiments may not be fully comparable with *in situ* flux measurements, these comparisons suggest that denitrification contributes significantly to biogenic nitric oxide production *in situ* in GBS. The much lower N₂O flux measured in SSW, which has ammonia oxidation and denitrification rates comparable to those in GBS (Dodsworth *et al.*, 2011), may be due to the relatively short water residence time of the SSW spring pool, preventing nitrate and nitrite from accumulating in sediment pore water or allowing N₂O to be removed primarily in aqueous form rather than by exchange with the atmosphere.

Enrichment, isolation, and identification of nitrate-reducing thermophiles

Heterotrophic micro-organisms isolated on four different field trips, using a variety of carbon substrates and four different enrichment strategies, were identified by 16S rRNA gene sequencing as members of the genera *Thermus*, *Anoxybacillus*, or *Geobacillus*. Only *T. thermophilus* (31 isolates) and *T. oshimai* (three isolates) were able to grow with nitrate as the terminal electron acceptor for anaerobic respiration (Table 2). *T. aquaticus* (four isolates) was not able to reduce nitrate, consistent with the description of that species (da Costa *et al.*, 2001). 16S rRNA gene fragments from Great Basin *T. thermophilus* isolates were 99.4–99.7% identical to the sequence from *T. thermophilus* HB8^T, differing by two conserved nucleotide substitutions from HB8, and possessing seven single-nucleotide polymorphisms, including a single-nucleotide insertion. Phylogenetic analysis of the *T. thermophilus* isolates showed no pattern with respect to isolation date or strategy (data not shown). *T. oshimai* isolates shared 99.5–99.6% identity with the sequence from *T. oshimai* SPS-17^T, differing by two conserved nucleotide substitutions from SPS-17^T and two single-nucleotide polymorphisms, including a single-nucleotide insertion.

Thermus thermophilus and *T. aquaticus* were each isolated from both GBS and SSW sediment and water, whereas *T. oshimai* was only isolated from SSW sediments. *T. thermophilus* and *T. aquaticus* were the dominant *Thermales* detected by 16S rRNA gene-based cultivation-independent censuses at GBS and SSW by both conventional clone library analysis (Costa *et al.*, 2009) and pyrosequencing analysis, where sequences representing close relatives of *T. thermophilus* comprised 0.13–2.89% of pyrotags obtained from GBS site B (Dodsworth *et al.*, 2011; J. Guy, J. Peacock, and B. Hedlund, unpublished results). In contrast, *T. oshimai* was not detected

in the cultivation-independent studies, despite sequencing of >550 16S rRNA gene clones and >100 000 16S rRNA gene pyrotags. Thus, both cultivation-dependent and cultivation-independent works suggest *T. oshimai* was less abundant in these spring samples and possibly implies a more important role of *T. thermophilus* in the N-cycle. Previously, quantitative PCR for *T. thermophilus narG* revealed sediment populations of $1.3\text{--}1.7 \times 10^6$ copies g⁻¹ sediment at the GBS B site in October 2008 (Dodsworth *et al.*, 2011) and 1.5×10^7 copies g⁻¹ sediment at the GBS A site in June 2009 (J. Dodsworth and B. Hedlund, unpublished data). Collectively these results suggest that close relatives of *T. thermophilus* can represent significant members of microbial communities in GBS and, because these isolates all produce N₂O as a terminal product of denitrification, that *Thermus* spp. may therefore contribute to the high observed N₂O flux.

Determination of nitrate reduction intermediates and final products

All nitrate-reducing *Thermus* isolates produced N₂O as the dominant product and failed to produce N₂ or ammonia within 72 h of cultivation with 9 mM nitrate as the sole terminal electron acceptor (data not shown). Selected isolates were grown for up to 120 h but still failed to produce N₂ or ammonia (Fig. 3; data not shown). In addition, mid-

exponential-phase cells grown with nitrate as the terminal electron acceptor were unable to produce gas or grow when transferred into the same medium with N₂O as the sole terminal electron acceptor.

To study the nitrate reduction phenotype in more detail, two isolates each from *Thermus* species were selected for further analysis based on their robust and consistent growth phenotype (Fig. 3). Consistent with the results described earlier, N₂O accumulated in the cultures over time, was the major end product of nitrate reduction, and was not reduced to N₂ during the course of the experiments. The conversion of nitrate to N₂O was nearly stoichiometric, with transient accumulation of nitrite. Production of N₂O commenced in mid- to late-exponential growth phase and continued during stationary phase, suggesting that denitrification can be, but is not necessarily, coupled to an increase in cell number in these strains. The complete removal of nitrate by *T. oshimai* strains suggests a higher affinity nitrate reductase or better maintenance of this activity during stationary phase than in *T. thermophilus*.

These *Thermus* isolates have a denitrification phenotype previously unknown in the *Thermales*. Other thermophiles that denitrify to N₂O as the final product have been described, including *Ferroplasma placidus* (Vorholt *et al.*, 1997), *Hydrogenivirga caldilitoralis* (Nakagawa *et al.*, 2004), *Petrobacter succinatimendens* (Salinas *et al.*, 2004), *Pseudoxanthomonas*




Fig. 3 Near stoichiometric conversion of NO₃⁻ to N₂O during growth. Cells were grown in *Thermus* medium under a He atmosphere and sampled periodically for quantification of cell density, NO₃⁻, and the possible reduction products NO₂⁻, N₂O, N₂, and NH₃. Increases in N₂ and NH₃ were never detected (data not shown). Bars for cell counts represent standard deviations. N species were measured once at each time point. Data are representative of two or more identical experiments for each strain.

taiwanensis (Chen *et al.*, 2002), and some strains of *Geobacillus stearothermophilus* (Gokce *et al.*, 1989). In addition, although *Pyrobaculum aerophilum* can reduce N₂O to N₂, nitrous oxide reductase activity is very low, so N₂O accumulates as the major product of denitrification (deVries & Schröder, 2002). Some other thermophiles can denitrify completely to N₂, including *Sulfurihydrogenibium subterraneum* (Takai *et al.*, 2003), *Hydrogenobacter thermophilus* (Suzuki *et al.*, 2001), some strains of *Geobacillus* (Mishima *et al.*, 2009), and some strains of *T. thermophilus* (Cava *et al.*, 2008); however, the kinetics of denitrification in these organisms has not been characterized, so the ratios of N₂O and N₂ produced are not known.

At present, it is difficult to ascertain whether incomplete denitrification is more common among thermophiles and hyperthermophiles than among mesophiles; however, the positive relationship between temperature and N₂O flux measured at GBS hint that incomplete denitrification may be more active at high temperatures. Interestingly, experiments performed at site B of GBS showed ~15- to 20-fold higher rates of N₂O production with an acetylene block, demonstrating that the physiological capacity for N₂O reduction is present (Dodsworth *et al.*, 2011). However, the acetylene block experiments were conducted in a closed system and thus may overestimate *in situ* rates of N₂O reduction. We postulate that in the natural spring environment, the continuous supply of electron donor and nitrate as an electron acceptor favors incomplete denitrification, with N₂O as the dominant end product. It is also possible that complete denitrification does occur under natural conditions through the action of yet-uncultivated complete denitrifiers in these sediments or through the use of released N₂O as a terminal electron acceptor by other organisms.

Analysis of narG fragments

Putative *narG* fragments were amplified from DNA extracts of all denitrifying *T. thermophilus* and *T. oshimai* strains tested. All *narG* fragments were most closely related to the *narG* of *T. thermophilus* HB8 with DNA identities of 96–98% for *T. thermophilus* and 93% for *T. oshimai* and deduced amino acid identities of 96–98% and 95%, respectively. Nucleic acid and deduced amino acid identities to other nitrate-reducing *Thermales*, *Meiothermus silvanus* DSM 9946 and *Oceanithermus profundus* DSM 14977, were 73–83% and 73–86%, respectively. The most closely related *narG* sequences outside the *Thermales* were in the *Deltaproteobacteria* genera *Anaeromyxobacter* and *Geobacter*, with <71% DNA identity and <65% amino acid identity to *Thermus narG*. *T. oshimai* JL-4 and *T. thermophilus* JL-18 each possessed one or more large plasmid, suggesting the possible conservation of the NCE megaplasmid in these organisms (data not shown). However, the close relationship between *narG* within each *Thermus* species, regardless of geographic location, suggests

that interspecies or intergenus transfer of the NCE does not occur. The apparent lack of N₂O reductase activity in both *T. thermophilus* and *T. oshimai* from the Great Boiling Spring system suggests a lack of selection for the maintenance of functional *nos* genes, the genetic basis for which will be a subject of future bioinformatic analysis of *T. thermophilus* and *T. oshimai* genomes.

CONCLUSION

The N₂O fluxes measured in the GBS source pool and high temperature shallow sediments (site B) are comparable to rates measured in a variety of aquatic habitats such as coastal marine environments (0–1142 μmol N₂O-N m⁻² day⁻¹ (Seitzinger, 1988; Corredor *et al.*, 1999; Chen *et al.*, 2010)), freshwater stream and riparian zone environments (~10–1214 μmol N₂O-N m⁻² day⁻¹ (Seitzinger, 1988; Heftig *et al.*, 2003)), and freshwater lakes [0–19.2 μmol N₂O-N m⁻² day⁻¹ (Seitzinger, 1988; Huttunen *et al.*, 2003)]. In light of the suggestion that more than 90% of N₂O emissions from river and estuary sediments are anthropogenic (Seitzinger *et al.*, 2000), the rates of N₂O flux from GBS are particularly high, as this geothermal system is not impacted by anthropogenic N. The high N₂O flux is, however, in general agreement with high rates of denitrification measured in GBS (Dodsworth *et al.*, 2011). These data suggest a model where the N-cycle in these springs is based on chemolithotrophic oxidation of source water ammonia by ammonia-oxidizing archaea, followed by respiration of nitrate or nitrite to gaseous products (denitrification) or back into ammonia (DNRA). The high flux of N₂O into the atmosphere from GBS may be due to the longer residence time of the main spring pool.

Thermus thermophilus and *T. oshimai* were the dominant nitrate-reducing thermophiles isolated in this study. All isolates produced N₂O as the major product of nitrate reduction, a denitrification phenotype not previously described in the *Thermales*, although known in other thermophiles. Incomplete denitrification to N₂O and respiratory nitrate ammonification may be more common among thermophiles than mesophiles, possibly spelling a different fate for nitrate and nitrite in high temperature geothermal environments. N₂O released by these organisms may be the source of the high N₂O flux from high temperature parts of GBS.

ACKNOWLEDGEMENTS

The authors thank David and Sandy Jamieson for access to GBS and SSW and Shannon Stockton and the hardworking students of Pyramid Lake Junior and Senior High School in Nixon, NV, for participation with enrichment strategy 3 in June 2009. We thank Kyle Costa for isolation and identification of strains PS1 and S1, and Jen Fisher, Jessica Newburn, and Jessica Guy for valuable discussions on drafts of the manuscript. This work was supported by NSF Grant Numbers

MCB-0546865, OISE-0968421, and EPS-9977809. A grant from the UNLV Research Infrastructure Award Program (RIAP) provided the GC-ECD and GC-TCD used for measurements of N₂O and N₂.

REFERENCES

- Afshar S, Kim C, Monbouquette HG, Schroder II (1998) Effect of tungstate on nitrate reduction by the hyperthermophilic archaeon *Pyrobaculum aerophilum*. *Applied and Environmental Microbiology* **64**, 3004–3008.
- Albuquerque L, Ferreira C, Tomaz D, Tiago I, Verissimo A, da Costa MS, Nobre MF (2009) *Meiothermus rufus* sp. nov., a new slightly thermophilic red-pigmented species and emended description of the genus *Meiothermus*. *Systematic and Applied Microbiology* **32**, 306–313.
- Albuquerque L, Rainey FA, Nobre MF, da Costa MS (2010) *Meiothermus granaticus* sp. nov., a new slightly thermophilic red-pigmented species from the Azores. *Systematic and Applied Microbiology* **33**, 243–246.
- Allen MB (1959) Studies with *Cyanidium caldarium*, an anonymously pigmented chlorophyte. *Archiv für Mikrobiologie* **32**, 270–277.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *Journal of Molecular Biology* **215**, 403–410.
- Bhandari B, Simlot MM (1986) Rapid micro-method for determination of nitrate in presence of nitrite for biochemical studies. *Indian Journal of Experimental Biology* **24**, 323–325.
- Blöchl E, Rachel R, Burggraf S, Hafenbradl D, Jannasch HW, Stetter KO (1997) *Pyrolobus fumarii*, gen. and sp. nov., represents a novel group of archaea, extending the upper temperature limit for life to 113 degrees C. *Extremophiles* **1**, 14–21.
- Burr MD, Botero LM, Young MJ, Inskeep WP, McDermott TR (2005) Observations concerning nitrogen cycling in a Yellowstone thermal soil environment. In *Geothermal Biology and Geochemistry in Yellowstone National Park* (eds Inskeep WP, McDermott TR), Montana State University Publications, Bozeman, MT, pp. 171–182.
- Castenholz RW (1969) Thermophilic blue-green algae and the thermal environment. *Bacteriological Reviews* **33**, 476–504.
- Cava F, Zafra O, da Costa MS, Berenguer J (2008) The role of the nitrate respiration element of *Thermus thermophilus* in the control and activity of the denitrification apparatus. *Environmental Microbiology* **10**, 522–533.
- Chen MY, Tsay SS, Chen KY, Shi YC, Lin YT, Lin GH (2002) *Pseudoxanthomonas taiwanensis* sp. nov., a novel thermophilic, N₂O-producing species isolated from hot springs. *International Journal of Systematic and Evolutionary Microbiology* **52**, 2155–2161.
- Chen GC, Tam NFY, Ye Y (2010) Summer fluxes of atmospheric greenhouse gases N₂O, CH₄, and CO₂ from mangrove soil in South China. *Science of the Total Environment* **408**, 2761–2767.
- Correador JE, Morell JM, Bauza J (1999) Atmospheric nitrous oxide fluxes from mangrove sediments. *Marine Pollution Bulletin* **38**, 473–478.
- Costa KC, Navarro JB, Shock EL, Zhang CL, Soukup D, Hedlund BP (2009) Microbiology and geochemistry of Great Boiling and Mud Hot Springs in the United States Great Basin. *Extremophiles* **13**, 447–459.
- da Costa MS, Nobre MF, Rainey F (2001) Genus *Thermus*. In *Bergey's Manual of Systematic Bacteriology* (eds Boone DR, Castenholz RW, Garrity GM), Springer, New York, 404–414.
- Dodsworth JA, Hungate B, de la Torre J, Jiang H, Hedlund BP (2010) Measuring nitrification, denitrification, and related biomarkers in continental geothermal ecosystems. *Methods in Enzymology* **486**, 171–203.
- Dodsworth JA, Hungate BA, Hedlund BP (2011) Ammonia oxidation, denitrification, and dissimilatory nitrate reduction to ammonium in two US Great Basin hot springs with abundant ammonia-oxidizing archaea. *Environmental Microbiology* **13**, 2371–2386.
- Eder W, Huber R (2002) New isolates and physiological properties of the Aquificales and description of *Thermocrinis albus* sp. nov. *Extremophiles* **6**, 309–318.
- Eder W, Ludwig W, Huber R (1999) Novel 16S rRNA gene sequences retrieved from highly saline brine sediments of Kebrut Deep, Red Sea. *Archives of Microbiology* **172**, 213–218.
- Eder W, Jahnke LL, Schmidt M, Huber R (2001) Microbial diversity of the brine-seawater interface of the Kebrut Deep, Red Sea, studied via 16S rRNA gene sequences and cultivation methods. *Applied and Environmental Microbiology* **67**, 3077–3085.
- EPA U.S. (2010) 2010 U.S. Greenhouse gas inventory report (EPA document #430-R-10-006). In (ed. EPA, U.S.).
- Gokce N, Hollocher TC, Bazylnski DA, Jannasch HW (1989) Thermophilic *Bacillus* sp. that shows the denitrification pathway of *Pseudomonas aeruginosa*. *Applied and Environmental Microbiology* **55**, 1023–1025.
- Gotz D, Banta A, Beveridge TJ, Rushdi AI, Simoneit BR, Reysenbach AL (2002) *Persephonella marina* gen. nov., sp. nov. and *Persephonella guaymasensis* sp. nov., two novel, thermophilic, hydrogen-oxidizing microaerophiles from deep-sea hydrothermal vents. *International Journal of Systematic and Evolutionary Microbiology* **52**, 1349–1359.
- Hafenbradl D, Keller M, Dirmeier R, Rachel R, Rosnagel P, Burggraf S, Huber H, Stetter KO (1996) *Ferroplasma placidum* gen. nov., sp. nov., a novel hyperthermophilic archaeum that oxidizes Fe²⁺ at neutral pH under anoxic conditions. *Archives of Microbiology* **166**, 308–314.
- Hall T (2005). BioEdit: biological sequence alignment editor for Win95/98/NT/2K/XP. URL <http://www.mbio.ncsu.edu/bioedit/bioedit.html>.
- Hart LT, Larson AD, McCleskey CS (1965) Denitrification by *Corynebacterium nephridii*. *Journal of Bacteriology* **89**, 1104–1108.
- Heftig MM, Bobbink R, de Caluwe H (2003) Nitrous oxide emission and denitrification in chronically nitrate-loaded riparian buffer zones. *Journal of Environmental Quality* **32**, 1194–1203.
- Huber R, Rosnagel P, Woese CR, Rachel R, Langworthy TA, Stetter KO (1996) Formation of ammonium from nitrate during chemolithoautotrophic growth of the extremely thermophilic bacterium *Ammonifex degensii* gen. nov. sp. nov. *Systematic and Applied Microbiology* **19**, 40–49.
- Huttunen JT, Alm J, Liikanen A, Juutinen S, Larmola T, Hammar T, Silvola J, Martikainen PJ (2003) Fluxes of methane, carbon dioxide and nitrous oxide in boreal lakes and potential anthropogenic effects on the aquatic greenhouse gas emissions. *Chemosphere-Global Change Science* **52**, 609–621.
- Johnson DB, Rolfe S, Hallberg KB, Iversen E (2001) Isolation and phylogenetic characterization of acidophilic microorganisms indigenous to acidic drainage waters at an abandoned Norwegian copper mine. *Environmental Microbiology* **3**, 630–637.
- MacIntyre S, Wanninkhof JP, Chanton PA (1995) Biogenic trace gases: measuring emissions from soil and water. In *Biogenic Trace Gases: Measuring Emissions from Soil and Water* (eds Matson PA, Harriss RC), Blackwell Science, New York, pp., 52–97.
- Matson PA, Naylor R, Ortiz-Monasterio I (1998) Integration of environmental, ergonomic, and economic aspects of fertilizer management. *Science* **280**, 112–115.

- Miroshnichenko ML, L'Haridon S, Jeanthon C, Antipov AN, Kostrikina NA, Tindall BJ, Schumann P, Spring S, Stackebrandt E, Bonch-Osmolovskaya EA (2003a) *Oceanithermus profundus* gen. nov., sp. nov., a thermophilic, microaerophilic, facultatively chemolithoheterotrophic bacterium from a deep-sea hydrothermal vent. *International Journal of Systematic and Evolutionary Microbiology* **53**, 747–752.
- Miroshnichenko ML, L'Haridon S, Necessian O, Antipov AN, Kostrikina NA, Tindall BJ, Schumann P, Spring S, Stackebrandt E, Bonch-Osmolovskaya EA, Jeanthon C (2003b) *Vulcanithermus mediatlanticus* gen. nov., sp. nov., a novel member of the family *Thermaceae* from a deep-sea hot vent. *International Journal of Systematic and Evolutionary Microbiology* **53**, 1143–1148.
- Mishima M, Iwata K, Nara K, Matsui T, Shigeno T, Omori T (2009) Cultivation characteristics of denitrification by thermophilic *Geobacillus* sp. strain TDN01. *The Journal of General and Applied Microbiology* **55**, 81–86.
- Nakagawa S, Nakamura S, Inagaki F, Takai K, Shirai N, Sako Y (2004) *Hydrogenivirga caldilitoris* gen. nov., sp. nov., a novel extremely thermophilic, hydrogen- and sulfur-oxidizing bacterium from a coastal hydrothermal field. *International Journal of Systematic and Evolutionary Microbiology* **54**, 2079–2084.
- Philippot L, Piutti S, Martin-Laurent F, Hallet S, Germon JC (2002) Molecular analysis of the nitrate-reducing community from unplanted and maize-planted soils. *Applied and Environmental Microbiology* **68**, 6121–6128.
- Pires AL, Albuquerque L, Tiago I, Nobre MF, Empadinhas N, Verissimo A, da Costa MS (2005) *Meiothermus timidus* sp. nov., a new slightly thermophilic yellow-pigmented species. *FEMS (Federation of European Microbiological Societies) Microbiology Letters* **245**, 39–45.
- Poli A, Romano I, Cordella P, Orlando P, Nicolaus B, Ceschi Berrini C (2009) *Anoxybacillus thermarum* sp. nov., a novel thermophilic bacterium isolated from thermal mud in Eugeanean hot springs, Abano Terme, Italy. *Extremophiles* **13**, 867–874.
- Ramirez S, Moreno R, Zafra O, Castan P, Valles C, Berenguer J (2000) Two nitrate/nitrite transporters are encoded within the mobilizable plasmid for nitrate respiration of *Thermus thermophilus* HB8. *Journal of Bacteriology* **182**, 2179–2183.
- Ramirez-Arcos S, Fernandez-Herrero LA, Berenguer J (1998) A thermophilic nitrate reductase is responsible for the strain specific anaerobic growth of *Thermus thermophilus* HB8. *Biochimica et Biophysica Acta* **1396**, 215–227.
- Romanek C, Zhang C, Mills G, Jones M, Paddock L (2005) Physiochemical heterogeneity in Nevada hot springs: implications for microbial diversity and function. *105th General Meeting of the American Society for Microbiology*.
- Salinas MB, Fardeau ML, Cayol JL, Casalat L, Patel BK, Thomas P, Garcia JL, Ollivier B (2004) *Petrobacter succinatimandens* gen. nov., sp. nov., a moderately thermophilic, nitrate-reducing bacterium isolated from an Australian oil well. *International Journal of Systematic and Evolutionary Microbiology* **54**, 645–649.
- Schlesinger WH (2009) On the fate of anthropogenic nitrogen. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 203–208.
- Seitzinger SP (1988) Denitrification in freshwater and coastal marine ecosystems: ecological and geochemical significance. *Limnology and Oceanography* **33**, 702–724.
- Seitzinger SP, Kroeze C, Styles RV (2000) Global distribution of N₂O emissions from aquatic systems: natural emissions and anthropogenic effect. *Chemosphere-Global Change Science* **2**, 267–279.
- Seitzinger S, Harrison JA, Böhlke JK, Bouwman AF, Lowrance R, Peterson B, Tobias C, Van Drecht G (2006) Denitrification across landscapes and waterscapes: a synthesis. *Ecological Applications* **16**, 2064–2090.
- Suzuki M, Cui ZJ, Ishii M, Igarashi Y (2001) Nitrate respiratory metabolism in an obligately autotrophic hydrogen-oxidizing bacterium, *Hydrogenobacter thermophilus* TK-6. *Archives of Microbiology* **175**, 75–78.
- Takai K, Kobayashi H, Nealson KH, Horikoshi K (2003) *Sulfurivibrio subterraneum* gen. nov., sp. nov., from a subsurface hot aquifer. *International Journal of Systematic and Evolutionary Microbiology* **53**, 823–827.
- Völkl P, Huber R, Drobner E, Rachel R, Burggraf S, Trincone A, Stetter KO (1993) *Pyrobaculum aerophilum* sp. nov., a novel nitrate-reducing hyperthermophilic archaeum. *Applied and Environmental Microbiology* **59**, 2918–2926.
- Vorholt JA, Hafenbradl D, Stetter KO, Thauer RK (1997) Pathways of autotrophic CO₂ fixation and of dissimilatory nitrate reduction to N₂O in *Ferroglobus placidus*. *Archives of Microbiology* **167**, 19–23.
- deVries S, Schröder I (2002) Comparison between the nitric oxide reductase family and its aerobic relatives, the cytochrome oxidases. *Biochemical Society Transactions* **30**, 662–667.
- Wilhelm E, Battino R, Wilcock RJ (1977) Low-pressure solubility of gases in liquid water. *Chemical Reviews* **77**, 219–262.
- Zafra O, Ramirez S, Castan P, Moreno R, Cava F, Valles C, Caro E, Berenguer J (2002) A cytochrome c encoded by the nar operon is required for the synthesis of active respiratory nitrate reductase in *Thermus thermophilus*. *FEBS Letters* **523**, 99–102.
- Zhang XQ, Zhang WJ, Wei BP, Xu BP, Zhu XF, Wu M (2010) *Meiothermus cateniformans* sp. nov., a slightly thermophilic species from north-eastern China. *International Journal of Systematic and Evolutionary Microbiology* **60**, 840–844.
- Zumft WG, Kroneck PM (2007) Respiratory transformation of nitrous oxide (N₂O) to dinitrogen by Bacteria and Archaea. *Advances in Microbial Physiology* **52**, 107–227.